


# Mathematics 1100Y – Calculus I: Calculus of one variable

TRENT UNIVERSITY, SUMMER 2011

## Solution to Assignment #10 Differential Dog Drag

Little human  $M$  is trying to walk big dog  $B$  in a backyard with a rectangular pool\*. With  $B$  keeping the  $3\text{ m}$  leash fully extended, they approach one corner of the pool. At the instant that  $B$  reaches the corner, the leash is extended in the direction of one of the sides, but then  $B$  spots squirrel  $S$  and runs off along the other side of the pool, dragging  $M$  along. At any given instant, the leash is fully extended and tangent to the curve that  $M$  is being dragged along.



Suppose we set up a Cartesian coordinate system so that the positive  $y$ -axis is on the edge of the pool that  $B$  runs off along, the origin is at the corner of the pool that  $B$  starts running from, and  $M$  is at  $(3, 0)$  when  $B$  starts running.


1. Find a function  $f(x)$  whose graph is the curve that  $M$  is dragged along, with the coordinate system set up as described above. [10]

HINT: If  $M$  is at  $(x, y)$  at some instant, where  $y = f(x)$ , the  $y$ -intercept of the tangent line always  $3\text{ m}$  from  $(x, y)$ . Recall, too, that the tangent line at  $(x, y)$  has slope  $m = \frac{dy}{dx} = f'(x)$ . Use all this to set up an equation involving  $\frac{dy}{dx}$  and then solve it for  $y$ .

SOLUTION. When  $M$  is at  $(x, f(x))$ , consider the right triangle whose hypotenuse is the leash, and hence has length  $3$ , and whose short sides are parallel to the axes. The base of this triangle, the side parallel to the  $x$ -axis, has length  $x - 0 = x$ ; let  $c$  be the length of the other short side, the side parallel to the  $y$ -axis.

---

\* The names and situation have been changed slightly to ~~protect the innocent~~ set this problem up.



By the Pythagorean Theorem,  $c^2 + x^2 = 3^2$ , so  $c = \sqrt{3^2 - x^2} = \sqrt{9 - x^2}$ . The slope of the leash – which is equal to  $\frac{dy}{dx}$  because the leash is tangent to the curve – is then  $\frac{\text{rise}}{\text{run}} = \frac{-c}{x} = -\frac{\sqrt{9 - x^2}}{x}$ . (Note that the slope must be negative because the leash goes down from left to right.) It follows that  $f'(x) = \frac{dy}{dx} = -\frac{\sqrt{9 - x^2}}{x}$ .

It remains to solve this equation for  $y = f(x)$ ; note that we also know from the initial setup that  $f(3) = 0$ .

*Attempt i.* One way to do the job is to use Maple. The worksheet-style command

```
> dsolve({diff(y(x),x)=-sqrt(9-x^2)/x,y(3)=0},y(x));
```

gives the result:

$$y(x) = -\sqrt{9 - x^2} + 3\text{arctanh}\left(\frac{3}{\sqrt{9 - x^2}}\right) + \frac{3}{2}I\pi$$

This solution, if you think about it, is a bit problematic: the  $I$  in the constant term represents the “imaginary” number  $i = \sqrt{-1}$ . You might ask yourself what it’s doing here, given since we’re supposed to be getting a real-valued function of the real variable  $x \dots$

*Attempt ii.* One can also do the job by hand using our knowledge of integration. It follows from the Fundamental Theorem of Calculus that  $f'(x) = \frac{dy}{dx} = -\frac{\sqrt{9 - x^2}}{x}$  and  $f(3) = 0$  imply that  $f(x) = \int_3^x -\frac{\sqrt{9 - t^2}}{t} dt$ . We will compute this integral using the trigonometric

substitution  $t = 3 \sin(\theta)$ , so  $dt = 3 \cos(\theta) d\theta$ :

$$\begin{aligned}
 f(x) &= \int_3^x -\frac{\sqrt{9-t^2}}{t} dt = -\int_{t=3}^{t=x} \frac{\sqrt{9-3^2 \sin^2(\theta)}}{3 \sin(\theta)} 3 \cos(\theta) d\theta \\
 &= \int_{t=x}^{t=3} \frac{3\sqrt{1-\sin^2(\theta)}}{\sin(\theta)} \cos(\theta) d\theta = 3 \int_{t=x}^{t=3} \frac{\sqrt{\cos^2(\theta)}}{\sin(\theta)} \cos(\theta) d\theta \\
 &= 3 \int_{t=x}^{t=3} \frac{\cos(\theta)}{\sin(\theta)} \cos(\theta) d\theta = 3 \int_{t=x}^{t=3} \frac{\cos^2(\theta)}{\sin(\theta)} d\theta = 3 \int_{t=x}^{t=3} \frac{1-\sin^2(\theta)}{\sin(\theta)} d\theta \\
 &= 3 \int_{t=x}^{t=3} \left( \frac{1}{\sin(\theta)} - \frac{\sin^2(\theta)}{\sin(\theta)} \right) d\theta = 3 \int_{t=x}^{t=3} (\csc(\theta) - \sin(\theta)) d\theta
 \end{aligned}$$

At this point we look up the antiderivative of  $\csc(x) \dots \ddot{\smile}$

$$\begin{aligned}
 &= 3 [-\ln(\csc(\theta) + \cot(\theta)) - (-\cos(\theta))] \Big|_{t=x}^{t=3} \\
 &= 3 \left[ \cos(\theta) - \ln \left( \frac{1}{\sin(\theta)} + \frac{\cos(\theta)}{\sin(\theta)} \right) \right] \Big|_{t=x}^{t=3}
 \end{aligned}$$

Note that  $\sin(\theta) = t/3$  and  $\cos(\theta) = \sqrt{1-\sin^2(\theta)} = \sqrt{1-t^2/9}$ .

$$\begin{aligned}
 &= 3 \left[ \sqrt{1-t^2/9} - \ln \left( \frac{1}{t/3} + \frac{\sqrt{1-t^2/9}}{t/3} \right) \right] \Big|_{t=x}^{t=3} \\
 &= \left[ 3\sqrt{1-t^2/9} - 3\ln \left( \frac{3}{t} + \frac{3}{t}\sqrt{1-t^2/9} \right) \right] \Big|_{t=x}^{t=3} \\
 &= \left[ \sqrt{9-t^2} - 3\ln \left( \frac{3}{t} + \frac{1}{t}\sqrt{9-t^2} \right) \right] \Big|_{t=x}^{t=3} \\
 &= \left[ \sqrt{9-t^2} - 3\ln \left( 3 + \sqrt{9-t^2} \right) - 3\ln \left( \frac{1}{t} \right) \right] \Big|_{t=x}^{t=3} \quad \text{But } \frac{1}{t} = t^{-1}, \text{ so } \dots \\
 &= \left[ \sqrt{9-t^2} - 3\ln \left( 3 + \sqrt{9-t^2} \right) - 3(-1)\ln(t) \right] \Big|_{t=x}^{t=3} \\
 &= \left[ \sqrt{9-3^2} - 3\ln \left( 3 + \sqrt{9-3^2} \right) + 3\ln(3) \right] \\
 &\quad - \left[ \sqrt{9-x^2} - 3\ln \left( 3 + \sqrt{9-x^2} \right) + 3\ln(x) \right] \\
 &= [0 - 3\ln(3+0) + 3\ln(3)] - \left[ \sqrt{9-x^2} - 3\ln \left( 3 + \sqrt{9-x^2} \right) + 3\ln(x) \right] \\
 &= 0 - \left[ -3\ln \left( 3 + \sqrt{9-x^2} \right) + 3\ln(x) + \sqrt{9-x^2} \right] \\
 &= 3\ln \left( 3 + \sqrt{9-x^2} \right) - 3\ln(x) - \sqrt{9-x^2}
 \end{aligned}$$

Whew! At least there are no imaginary terms ...  $\square$

NOTE: The curve that occurs in this problem is called a *tractrix*.