

# Mathematics 1110H – Calculus I: Limits, Derivatives, and Integrals

TRENT UNIVERSITY, Fall 2023

## Assignment #3

### The Limit Game and a Differential Equation

*Due\* just before midnight on Friday, 6 October.*

The usual  $\varepsilon - \delta$  definition of limits,

DEFINITION.  $\lim_{x \rightarrow a} f(x) = L$  exactly when for every  $\varepsilon > 0$  there is a  $\delta > 0$  such that for any  $x$  with  $|x - a| < \delta$  we are guaranteed to have  $|f(x) - L| < \varepsilon$  as well.

is pretty hard to wrap your head around the first time or three for most people. Here is less common version of the definition, equivalent to the standard one, which recasts the confusing logical structure of the standard definition in terms of a game:

ALTERNATE DEFINITION. The *limit game* for  $f(x)$  at  $x = a$  with target  $L$  is a three-move game played between two players  $A$  and  $B$  as follows:

1.  $A$  moves first, picking a small number  $\varepsilon > 0$ .
2.  $B$  moves second, picking another small number  $\delta > 0$ .
3.  $A$  moves third, picking an  $x$  that is within  $\delta$  of  $a$ , *i.e.*  $a - \delta < x < a + \delta$ .

To determine the winner, we evaluate  $f(x)$ . If it is within  $\varepsilon$  of the target  $L$ , *i.e.*  $L - \varepsilon < f(x) < L + \varepsilon$ , then player  $B$  wins; if not, then player  $A$  wins.

With this idea in hand,  $\lim_{x \rightarrow a} f(x) = L$  means that player  $B$  has a winning strategy in the limit game for  $f(x)$  at  $x = a$  with target  $L$ ; that is, if  $B$  plays it right,  $B$  will win no matter what  $A$  tries to do. (Within the rules ... :-)  
Conversely,  $\lim_{x \rightarrow a} f(x) \neq L$  means that player  $A$  is the one with a winning strategy in the limit game for  $f(x)$  at  $x = a$  with target  $L$ .

The game definition of limits isn't really better or worse than the usual  $\varepsilon - \delta$  definition, but each is easier for some people to understand, and the exercise in trying it both ways usually helps in understanding what is really going on with limits.

1. Use the alternate definition of limits to verify that  $\lim_{x \rightarrow 3} (-3x + 4) = -5$ . [2.5]

*Hint:* Try using the usual  $\varepsilon - \delta$  definition of limits first to work out a winning strategy for player  $B$ . Note that player  $B$  only gets one move: picking a  $\delta$  after player  $A$  plays an  $\varepsilon$ .

2. Use the alternate definition of limits to verify that  $\lim_{x \rightarrow -1} (2x - 3) \neq 0$ . [2.5]

*Hint:* You need a winning strategy for player  $A$ . What is a suitable  $\varepsilon$  to play on the first move that will defeat any  $\delta$  that player  $B$  might choose?

*More on page 2!*

---

\* You should submit your solutions via Blackboard's Assignments module, preferably as a single pdf. If submission via Blackboard fails, please submit your work to your instructor by email or on paper.

If you want to get started on this part of the assignment before attending your lab in MATH 1110H, skimming and later referring to as necessary to Sections 4.22.1 and 4.22.2 of Gregory Bard's book *Sage for Undergraduates* (in the SageMath folder in the Course Content section on Blackboard) is probably going to be useful.

- 3.** Use SageMath to solve for  $y$  as a function of  $x$  if  $\frac{dy}{dx} = -2y$  and it is also required that  $y = 1$  when  $x = 0$ . [4]

*Hint:* This is a job for the `desolve` command.

- 4.** Verify by hand that the solution you obtained using SageMath in question **3** is indeed a solution to the given differential equation,  $\frac{dy}{dx} = -2y$ , with the given initial condition,  $y = 1$  when  $x = 0$ . [1]