

# Mathematics 1550H – Introduction to probability

TRENT UNIVERSITY, Summer 2016

## Some Common Distributions

*The Short Form*

### Discrete

1. *Discrete Uniform.*  $n$  equally likely outcomes for some  $n \geq 1$ .

*Probability function:*  $m(\omega) = \frac{1}{n}$ .

Expected value and variance of a random variable  $X$  on  $\Omega$  depend on just what values  $X$  assigns to each outcome  $\omega \in \Omega$ .

2. *Bernoulli Trial.* Two outcomes with probability of success  $p$  and of failure  $q = 1 - p$ .  $X$  counts successes.

*Probability function:*  $m(1) = P(\text{success}) = p$  and  $m(0) = P(\text{failure}) = q$ .

*Expected value:*  $\mu = E(X) = p$       *Variance:*  $\sigma^2 = V(X) = pq$

3. *Binomial.*  $n$  Bernoulli trials, with probability of success  $p$  and of failure  $q = 1 - p$ .  $X$  counts successes.

*Probability function:*  $m(k) = P(k \text{ successes}) = \binom{n}{k} p^k q^{n-k}$ , where  $0 \leq k \leq n$ .

*Expected value:*  $\mu = E(X) = np$       *Variance:*  $\sigma^2 = V(X) = npq$

4. *Geometric.* Bernoulli trials repeated until the first success, with probability of success  $p$  and of failure  $q = 1 - p$ .  $X$  counts the number of trials required.

*Probability function:*  $m(k) = P(\text{first success on } k\text{th trial}) = q^{k-1} p$ , where  $k \geq 1$ .

*Expected value:*  $\mu = E(X) = \frac{1}{p}$       *Variance:*  $\sigma^2 = V(X) = \frac{q}{p^2}$

5. *Negative Binomial.* Bernoulli trials repeated until the  $k$ th success, with probability of success  $p$  and of failure  $q = 1 - p$ .  $X$  counts the number of trials required.

*Probability function:*  $m(x) = P(k \text{ success on } x\text{th trial}) = \binom{x-1}{k-1} p^k q^{x-k}$

*Expected value:*  $\mu = E(X) = \frac{k}{p}$       *Variance:*  $\sigma^2 = V(X) = \frac{kq}{p^2}$

6. *Poisson.* Used to approximate the Binomial distribution when  $n$  is large.

*Probability function:*  $m(x) = P(x) = \frac{e^{-\lambda} \lambda^x}{x!}$

*Expected value:*  $\mu = E(X) = \lambda$       *Variance:*  $\sigma^2 = V(X) = \lambda$

### Continuous

7. *Continuous Uniform.*

*Density function:*  $f(t) = \begin{cases} \frac{1}{b-a} & a \leq t \leq b \\ 0 & \text{otherwise} \end{cases}$

*Expected value:*  $\mu = E(X) = \frac{a+b}{2}$       *Variance:*  $\sigma^2 = V(X) = \frac{(b-a)^2}{12}$

8. Exponential.

$$\text{Density function: } f(t) = \begin{cases} \lambda e^{-\lambda t} & t \geq 0 \\ 0 & t < 0 \end{cases}$$

$$\text{Expected value: } \mu = E(X) = \frac{1}{\lambda} \quad \text{Variance: } \sigma^2 = V(X) = \frac{1}{\lambda^2}$$

9. Standard normal.

$$\text{Density function: } \varphi(t) = \frac{1}{\sqrt{2\pi}} e^{-t^2/2}$$

$$\text{Expected value: } \mu = E(X) = 0 \quad \text{Variance: } \sigma^2 = V(X) = 1$$

10. Normal. . . . with mean  $\mu$  and standard deviation  $\sigma$ .

$$\text{Density function: } f(t) = \frac{1}{\sigma\sqrt{2\pi}} e^{-(t-\mu)^2/2\sigma^2}$$

$$\text{Expected value: } E(X) = \mu \quad \text{Variance: } V(X) = \sigma^2$$