

Mathematics 3260H – Geometry II: Projective and non-Euclidean geometry  
TRENT UNIVERSITY, Winter 2015

Assignment #13 = 8 + 5\*

Free Completion

Due on Thursday, 5 March, 2015.

Suppose  $\mathcal{C}_0 = (\mathcal{P}_0, \mathcal{L}_0, I_0)$  is a configuration consisting of a set of points,  $\mathcal{P}_0$ , a set of lines,  $\mathcal{L}_0$ , and a relation of incidence,  $I_0$ , specifying which points are incident with which lines. For each  $n \geq 0$ , given the configuration  $\mathcal{C}_n = (\mathcal{P}_n, \mathcal{L}_n, I_n)$ , define the configuration  $\mathcal{C}_{n+1} = (\mathcal{P}_{n+1}, \mathcal{L}_{n+1}, I_{n+1})$  as follows:

- $\mathcal{P}_{n+1}$  includes all the points of  $\mathcal{P}_n$ , together with a new and distinct point of intersection for every pair of lines in  $\mathcal{L}_n$  which do not already have a common point of intersection in  $\mathcal{C}_n$ .
- $\mathcal{L}_{n+1}$  includes all the lines of  $\mathcal{L}_n$ , together with a new and distinct line joining each pair of points in  $\mathcal{P}_n$  which do not already have a line joining them in  $\mathcal{C}_n$ .
- $I_{n+1}$  is  $I_n$ , plus the incidences involving the added points and lines described above.

Finally, let  $\mathcal{P} = \bigcup_{n=0}^{\infty} \mathcal{P}_n$ ,  $\mathcal{L} = \bigcup_{n=0}^{\infty} \mathcal{L}_n$ , and  $I = \bigcup_{n=0}^{\infty} I_n$ . The incidence structure  $\mathcal{C} = (\mathcal{P}, \mathcal{L}, I)$  is the *free completion* of the configuration  $\mathcal{C}_0 = (\mathcal{P}_0, \mathcal{L}_0, I_0)$ .

1. Show that the free completion of an arbitrary (non-empty) configuration must satisfy axioms **I** and **II** for a projective plane. [5]
2. Give an example of a (non-empty) configuration whose free completion is *not* a projective plane. [1]
3. Assume  $\mathcal{C}_0$  consists of four points and no lines. Show that the free completion  $\mathcal{C}$  of  $\mathcal{C}_0$  is a projective plane. [4]

---

\* Toby ...